

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 June 2002 (13.06.2002)

PCT

(10) International Publication Number
WO 02/45695 A2

(51) International Patent Classification⁷: **A61K 9/22**

(21) International Application Number: PCT/CA01/01721

(22) International Filing Date: 4 December 2001 (04.12.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/251,751 5 December 2000 (05.12.2000) US

(71) Applicant and

(72) Inventor: MACGREGOR, Alexander [CA/CA]; 34
White Avenue, Scarborough, Ontario M1C 1P2 (CA).

(74) Agents: SECHLEY, Konrad, A. et al.; Gowling Lafleur
Henderson LLP, Suite 2600, 160 Elgin Street, Ottawa, On-
tario K1P 1C3 (CA).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA,
ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR,
GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent
(BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR,
NE, SN, TD, TG).


Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: HYDROSTATIC DELIVERY SYSTEM FOR CONTROLLED DELIVERY OF AGENT

Dissolution Profile of RANITIDINE HCL 60 mg Extended Release Tablet



(57) Abstract: The present invention provides a hydrostatic delivery system comprising a hydrostatic couple and an agent of interest. The hydrostatic couple comprises, at least one hydrodynamic fluid-imbibing polymer, and at least one hydrostatic pressure modulating agent. This delivery system has the ability to control the release of one or more agent of interest within a fluid environment following zero-order kinetics.



WO 02/45695 A2

- 1 -

HYDROSTATIC DELIVERY SYSTEM FOR CONTROLLED DELIVERY OF AGENT

5 The present invention relates to a delivery system for the controlled release of an agent of interest, as well as compositions and methods of preparation of the delivery system. More particularly, the present invention provides a hydrostatic pressure-activated delivery system for dispensing an agent of interest to an environment of use.

BACKGROUND OF THE INVENTION

10

The clinical advantages of controlled and patterned delivery of therapeutic agents are well established in the art. Many of the desirable attributes of controlled release pharmaceutical preparations stem from their ability to deliver predetermined quantities of one or more active agent(s) with a high degree of precision over a desired time frame.

15

Delivery devices and systems for the controlled release of active agents are generally characterized as either diffusion controlled delivery systems; erosion controlled systems, osmotic dispensing devices, or combinations of diffusion and erosion control. These devices and systems are derived from various compositions and techniques such as matrix processes, core embedding processes, coating processes as well as osmotically activated processes. Exemplifying these delivery devices are a broad range of systems from time release capsules whose contents have coatings that erode at different rates, diffusion-controlled matrix tablets with hydro-swellable barriers, and controlled release-rate tablets which operate by osmosis. Irrespective of the mechanism underlying the controlled release of an agent of interest, it is desired that a delivery system be characterized by a constant and reproducible *in-vivo* pharmacokinetic response facilitated by zero-order release kinetics (i.e. where the release of an agent of interest, for example a pharmaceutical agent, is independent of its own concentration).

20
25
30

U.S. 4,601,894, 4,687,757, 4,680,323, 4,994,276 disclose controlled release delivery devices based on matrix systems. These matrix systems are generally known to

- 2 -

lack the ability to release pharmaceutical agents according to zero-order kinetics. (e.g. S.D. Bruck, *Controlled Drug Delivery*, Vol. I and II, CRC Press (1983)).

5 Core embedding or core coated delivery systems have been disclosed, for example in U.S.3,538,214. This document describes a diffusion-controlled device in which a tablet core containing the active ingredient, is surrounded by a water insoluble coating. The insoluble film coating has been modified with modifying agents that are soluble to the external fluids in the gastrointestinal tract.

10 U.S. 3,845,770 and 3,916,899 disclose osmotic devices comprising a core composition of an active agent in combination with an osmotically effective solute, that is enclosed by an insoluble semi-permeable wall having a release capacity. The release characteristics of these devices have been improved through modifications disclosed, for example, in U.S. 4,624,847, 5,082,668. In principle, osmotic delivery employs one or
15 more osmotic pressure adjuvants, for example a salt, and one or more components involved in expansion, for example a polymer, to deliver an agent of interest to a fluid environment over a period of time. The osmotic pressure adjuvants present in the delivery device are used to cause the influx of water by osmosis, through a semi-permeable wall, while the component involved with expansion absorbs liquid, expands, and acts to drive
20 out the agent of interest from the interior of the osmotic device in a controlled and constant manner. Such systems are capable of zero-order release kinetics.

25 A disadvantage with coated delivery systems as well as osmotic devices, is that any damage to the wall or shell results in the premature release of the pharmaceutical agent within a short period of time causing what is known in the art as "dose dumping". Patient safety is jeopardized as a result of side effects and possible toxicity from high levels of an agent of interest, for example a pharmacological agent, being released within the blood stream over a short period of time.

30 While attempts have been made to minimize the safety risks associated with conventional single unit delivery devices by developing multiple unit osmotic pumps, these embellishments have led to increased manufacturing costs (e.g. S.D. Bruck, *supra*).

- 3 -

Similarly, osmotic delivery systems typically comprise one or more openings for the passage of an agent from the delivery device to the environment. The manufacture of the openings within the delivery device may be carried out using one or more laser drills (e.g. U.S. 3,845,770 and 3,916,899). The additional steps and machines required for the manufacture of fixed openings within the delivery device also increase the cost of
5 manufacture of such delivery devices.

While delivery systems have been designed that reduce the risk levels to the patient, there still remains significant and inherent shortcomings in osmotic devices, in part due to their reliance on the need for an osmotic gradient to be established between
10 the contents of the device and the fluid environment as well as the need for fixed opening(s) for the delivery of the agent. A blockage of the opening(s) either during storage or handling prior to patient consumption or due to the imminent interaction with dietary contents such as solid food particulate, or simply due to adherence to the
15 gastrointestinal cell wall, will alter the osmotic gradient and severely impair the performance of the osmotic device.

In addition, fluctuating osmolarity in the environment of use, such as the human gastro-intestinal tract, impacts on the reproducibility and performance of osmosis-
20 dependent devices. It is well known that the osmolarity of human gastro-intestinal fluid is imminently variable in the fed and fasted states. There can be a substantial increase of up to two fold in the fed state within the individual (J.B. Dressman, *Physiological Aspects of the Design of Dissolution Tests*, Scientific Foundation for Regulating Drug Product Quality- AAPS Press 1997). These natural variables are further pronounced by
25 diets containing varying salt and electrolyte contents. The performance of osmotically driven delivery devices is dependant upon many physiological variables and the dietary habits of patients. For example, side effects within patients (the "flame-cutter effect") arising from the concentrated release of a pharmaceutical agent from the release opening(s) of osmotic systems has led to the withdrawal of preparations comprising
30 Indomethacin.

- 4 -

Additionally, some active agents possess chemical properties that are comparable in ionic strengths to those of strong electrolytes and salts commonly used as osmotic adjuvants. In these instances, and due to different pH environments in the gastrointestinal tract, agents comprising significant ionic strength will manifest varying degrees of ionization that may compromise the predictable performance of the osmotic device. Osmotically active therapeutic agents with ionic strengths comparable to that of osmotic adjuvants, and that are localized within osmotically driven devices, will act as osmotic agents and enhance the osmotic influx of water from the fluid environment. Similarly, agents having high ionic strength may also cause variations in the osmolarity of the adjacent fluid environment upon their release from the delivery device. Therefore, osmotically-driven devices comprising agents characterised as having a high ionic strength, lack self-regulation.

A delivery system that is not readily influenced by minor changes to its physical form, intrinsic properties of an active agent (e.g. ionic strength), or variables in the environment of use (e.g. varying osmolarity of the human gastrointestinal tract and factors such as the dietary contents), can be reliably programmed to deliver the agent in a pre-determined manner with increased accuracy and precision. therefore, there remains within the art a need for a reliable zero-order drug delivery system, where the release of an agent is independent of its own concentration, that provides controlled drug delivery of an active agent to an environment of use and that is independent of physiological variables of the environment of use, as well as the intrinsic properties of the active agent.

It is an object of the invention to overcome disadvantages of the prior art.

The above object is met by the combinations of features of the main claims, the sub-claims disclose further advantageous embodiments of the invention.

- 5 -

SUMMARY OF THE INVENTION

The present invention relates to a delivery system for the controlled release of an agent of interest, as well as compositions and methods of preparation of the delivery system. More particularly, the present invention provides a hydrostatic pressure-activated delivery system for dispensing an agent of interest to an environment of use.

According to the present invention there is provided a hydrostatic delivery system comprising a hydrostatic couple and an agent of interest.

10

The present invention pertains to a hydrostatic delivery system comprising a hydrostatic couple comprising at least one hydrodynamic fluid-imbibing polymer, and at least one hydrostatic pressure modulating agent. Preferably, the hydrodynamic fluid-imbibing polymer is a cross-linked polymer with a swelling capacity in a fluid environment of between about 1 weight% to about 3000 weight%. Preferably, the cross-linked polymer is present from about 4 weight% to about 96 weight% of the total formulation. Also, it is preferred that the hydrostatic pressure modulating agent is a cross-linked, rapidly swelling polymer with a swelling capacity in a fluid environment of between about 0.5 weight% to about 500 weight%. Preferably the cross-linked, rapidly swelling polymer is present from about 1 weight% to about 50 weight% of the total formulation.

15
20

This invention further embraces a hydrostatic delivery system as defined above, wherein the hydrodynamic polymer and the hydrostatic pressure modulating agent are present at a ratio from about 99:1 to about 50:50 by weight.

25

The present invention also provides for the hydrostatic delivery system as defined above wherein the hydrostatic pressure modulating agent further comprises an expansion source, selected from the group consisting of a carbon-dioxide precursor, an oxygen precursor, and a chlorine dioxide precursor. Preferably, when the hydrodynamic polymer comprises a carbon dioxide precursor, oxygen precursor or chlorine dioxide precursor,

30

- 6 -

the hydrodynamic polymer and the hydrostatic pressure modulating agent are present in a ratio from about 99:1 to about 70:30 by weight.

5 According to the present invention, a hydrostatic delivery system is provided as defined above, comprising a hydrodynamic fluid-imbibing polymer selected from the group consisting of:

- i) an acrylic-acid polymer cross-linked with allylsucrose or allylpentaerythritol;
- ii) one or more starch derivatives cross-linked by Epichlorhydrin, Phosphorous oxychloride (POCl_3), or Sodium trimetaphosphate;
- 10 iii) a polyglucan;
- iv) a crosslinked polyacrylate resin;
- v) a crosslinked polyethylenimine;
- vi) a crosslinked polyallylamine, and combinations thereof, and

15 a hydrostatic pressure modulating agent selected from the group consisting of:

- i) homopolymers of cross-linked N-vinyl-2-pyrrolidone;
- ii) a rapidly expanding cross-linked cellulose derivative; and combinations thereof.

20 The present invention also provides for the hydrostatic delivery system as defined above, wherein the dosage form is a multiparticulate matrix tablet, or capsule. The hydrostatic delivery system may also comprising an enteric coating or one or more pH sensitive barrier polymers. The hydrostatic delivery system may be:

- i) a matrix-type solid compact, made by a compression or pelletization, a matrix-type extrusion spheroid, made by a wet or dry extrusion;
- 25 ii) be granulated or microencapsulated to form particulates that may be compressed into solid compacts or filled into capsules; or
- iii) spheroidal, compact, comprising dry blends, filled into capsules or suspended in a suitable liquid vehicle.

30

The present invention also embraces the hydrostatic delivery system as defined above, wherein the agent of interest is selected from the group consisting of analgesic,

- 7 -

anti-inflammatory, antimicrobial, amoebicidal, trichomonocidal agents, anti-parkinson, anti-malarial, anticonvulsant, anti-depressants, antiarthritics, anti-fungal, antihypertensive, antipyretic, anti-parasite, antihistamine, alpha-adrenargic agonist, alpha blocker, anesthetic, bronchial dilator, biocide, bactericide, bacteriostat, beta adrenergic
5 blocker, calcium channel blocker, cardiovascular drug, contraceptive, decongestants, diuretic, depressant, diagnostic, electrolyte, hypnotic, hormone, hyperglycemic, muscle relaxant, muscle contractant, ophthalmic, parasympathomimetic, psychic energizer, sedative, sympathomimetic, tranquilizer, urinary, vaginal, viricide, vitamin, non-steroidal anti-inflammatory, angiotensin converting enzyme inhibitors, polypeptide, proteins, and
10 sleep inducers.

This summary of the invention does not necessarily describe all necessary features of the invention but that the invention may also reside in a sub-combination of the described features.

15

- 8 -

BRIEF DESCRIPTION OF THE DRAWINGS

These and other features of the invention will become more apparent from the following description in which reference is made to the appended drawings wherein:

5 **FIGURE 1** shows the change in dynamic profile of a prior art delivery system.

FIGURE 2 shows the change in the dynamic profile of one formulation of a hydrostatic delivery system of the present invention.

10 **FIGURE 3** shows the dissolution profile of the release of an agent of interest from the hydrostatic delivery system of the prior art exhibiting non-zero (exponential) kinetics.

15 **FIGURE 4** shows the dissolution profile of the release of an agent of interest from the hydrostatic delivery system of the present invention exhibiting zero-order kinetics.

20 **FIGURE 5** shows the dissolution profile of the release of an agent of interest, Rantidine hydrochloride (60mg) from the hydrostatic delivery system of the present invention exhibiting zero-order kinetics.

25 **FIGURE 6** shows the dissolution profile of the release of an agent of interest, Tramadol hydrochloride (200mg) from the hydrostatic delivery system of the present invention exhibiting zero-order kinetics.

- 9 -

DESCRIPTION OF PREFERRED EMBODIMENT

5 The present invention relates to a delivery system for the controlled release of an agent of interest, as well as compositions and methods of preparation of the delivery system. More particularly, the present invention provides a hydrostatic pressure-activated delivery system for dispensing an agent of interest to an environment of use.

10 The following description is of a preferred embodiment by way of example only and without limitation to the combination of features necessary for carrying the invention into effect.

As used herein, the term "hydrodynamic fluid-imbibing polymer" means any polymer, synthetic or otherwise, that absorbs water or any fluid composition, in part by capillarity with a corresponding dynamic increase in volume and mass. By "capillarity" it is meant the passage of solvent into a solid polymer, for example a hydrodynamic fluid-imbibing polymer, as a result of differential pressure within the pore structure of the polymer and the fluid environment. Capillary uptake by the polymer is initiated by wetting and is dependent on the surface tension of the fluid and structural composition of the polymer.

15
20

As described in more detail below, and without wishing to be bound by theory, when two different intimately mixed fluid-imbibing polymers are compacted and exposed to a fluid environment such as water or biological fluid, the mixture of fluid imbibing polymers will absorb the fluid according to the individual contribution and propensity for water imbibition of the component polymers. If the rates and extents of fluid absorption are substantially different for the two polymers, and provided that the rate of volume expansion is greater for the polymer in lower concentration within the mixture, a differential dynamic volume expansion results. Such differential expansion essentially creates an internal stress that will ordinarily disrupt the compact causing complete disintegration. If the polymer present in larger molar concentration within the mixture is cross-linked thereby creating a microporous structure, the microporous structure of the

25
30

- 10 -

polymer network offers a resistance to the internal stress from the other rapidly expanding polymer. This resistance creates a positive hydrostatic differential pressure that increases the efflux of imbibed fluid to the external media. Under these conditions the volume of the mixture of the fluid-imbibing polymers continues to increase at a diminishing rate
5 determined by the hydrostatic pressure driven fluid efflux. At a given point the rates of volume increase will equal the rate of volume efflux and a dynamic steady state develops whereby the volume change is negligible. The dynamic steady state proffers a constant volume and surface area for both fluid influx and efflux. The delivery system operating on these principles is capable of controlling the release of both soluble and poorly soluble
10 agents of interest by effectively changing the rate of volume efflux through hydrostatic pressure modulation. The hydrostatic pressure of the delivery system may be increased by adding an expansion source, for example a mixture of alkaline and acidic agent as described below.

15 As used herein, "hydrostatic delivery system" refers to a composition that controls the release of an agent of interest contained therein, using non-osmotic hydrostatic differential pressure.

By "agent of interest" and "beneficial agent" it is meant one or more compounds
20 or mixture of compounds that can be released from the delivery system of the present invention to produce a desired or beneficial result. The agent of interest can be soluble in the fluid that is imbibed by the delivery system or it can have limited solubility in the imbibed fluid and be mixed with an effective solubilizer to enhance its solubility or a suitable excipient to retard its solubility. The agent of interest can be in the delivery
25 system in form of solid particles, granules, microencapsulated solid, microencapsulated liquid, powder and coated particles, for example, the agent of interest may comprise a plurality of discrete active particulates. Water insoluble agents of interest can be used in form that renders it water soluble and upon release from the delivery system, is converted to its original, or biologically active form, by enzyme hydrolysis, by pH, or a metabolic
30 processes depending on the environment of use.

- 11 -

Examples of beneficial agents are disclosed in Remington's Pharmaceutical Sciences (16th Ed., 1980, published by Mack Publishing Co., Easton, Pa.; and in The Pharmacological Basis of Therapeutics, by Goodman and Gilman, 6th Ed., 1980, published by The MacMillan Company, London). Furthermore, an agent of interest may

5 be selected from the following compounds, however, it is to be understood that the following compounds is not meant to be exhaustive. Many other agents of interest will certainly work in the hydrostatic delivery system of this invention. For example, agents of interest include, but are not limited to, pesticides, herbicides, germicides, biocides, fungicides, algicides, insecticides, rodenticides, antioxidants, preservatives, plant growth

10 inhibitors, plant growth promoters, chemical reactants, disinfectants, sterilization agents, foods, fermentation agents, food supplements, cosmetics, nutrients, vitamins, pharmaceutical drugs, nutraceuticals, vitamins, sex sterilants, fertility promoters, fertility inhibitors, microorganism attenuators, air purifiers, or other agents that benefit the environment of their use. By "drug", it is meant any therapeutically or pharmacologically

15 active substances that produce a localized or systemic effect or effects in animals, for example, but not limited to mammals, humans and primates. The expression "drug formulation" as used herein means the drug, by itself or the drug along with other excipients in an intimate mixture with a hydrostatic couple as described herein.

20 Therapeutic or pharmacologically active substances also include, but are not limited to, analgesic, anti-inflammatory, antimicrobial, amoebicidal, trichomonocidal agents, anti-parkinson, anti-malarial, anticonvulsant, anti-depressants, antiarthritics, anti-fungal, antihypertensive, antipyretic, anti-parasite, antihistamine, alpha-adrenargic agonist, alpha blocker, anesthetic, bronchial dilator, biocide, bactericide, bacteriostat,

25 beta adrenergic blocker, calcium channel blocker, cardiovascular drug, contraceptive, decongestants, diuretic, depressant, diagnostic, electrolyte, hypnotic, hormone, hyperglycemic, muscle relaxant, muscle contractant, ophthalmic, parasympathomimetic, psychic energizer, sedative, sympathomimetic, tranquilizer, urinary, vaginal, viricide, vitamin, non-steroidal anti-inflammatory, angiotensin converting enzyme inhibitors,

30 polypeptide, proteins, sleep inducers.

- 12 -

Other agents of interest include, but are not limited to, organic and inorganic compounds in various forms, such as charged molecules, molecular complexes, pharmacologically acceptable salts such as hydrochlorides, hydrobromides, palmitate, phosphate, sulphate laurylate, nitrate, borate, maleate, tartrate, acetate, salicylate and
5 oleate. Prodrugs and derivatives of drugs such as esters, ethers and amides.

An agent of interest also includes drugs that act on the peripheral nerves, for example, but not limited to, adrenergic receptors, cholinergic receptors, nervous system, skeletal muscles, cardiovascular, smooth muscles, blood circulatory system, synaptic
10 sites, neuroeffector junctional sites, endocrine and hormone systems, immunological system, reproductive system, skeletal system, autocoid systems, alimentary and excretory systems, inhibitory of autocoids and histamine systems, those materials that act on the central nervous system such as hypnotics and sedatives, including pentobarbital sodium, phenobarbital, secobarbital, thiopental and mixtures thereof; heterocyclic hypnotics such
15 as dioxopiperidines and glutarimides; hypnotics and sedatives such as amides and ureas, exemplified by diethylisovaleramide and .alpha.-bromoisovaleryl urea; hypnotic and sedative urethanes and disulfanes; psychic energizers such as isocoboxazid, nialamide, phenelzine, imipramine, amitriptyline hydrochloride, tranlycypromine and pargylene; and protryptiline hydrochloride, tranquilizers such as chlorpromazine, promazine,
20 fluphenzaine, reserpine, deserpidine, meprobamate, and benzodiazepines such as chlordiazepoxide; anticonvulsants such as primidone, enitabas, diphenylhydantion, ethyltion, pheneturide and ethosuximide; muscle relaxants and antiparkinson agents such as mephenesin, methocarbomal, cyclobenzaprine trihexylphenidyl, levodopa/carbidopa, and biperiden; antihypertensives such as .alpha.-methyldopa and L-.beta.-3-4-
25 dihydroxyphenylalanine, and pivaloyloxyethyl ester of .alpha.-methyldopa hydrochloride dihydrate; analgesics such as morphine, codeine, meperidine, nalorphine; antipyretics and anti-inflammatory agents such as aspirin, indomethacin, sodium indomethacin trihydrate salicylamide, naproxen, colchicine, fenoprofen, sulindac, diflunisal, diclofenac, indoprofen and sodium salicyl-amide; local anesthetics such as procaine, lidocaine,
30 maepaine, piperocaine, tetracaine and dibucane; antispasmodics and muscle contractants such as atropine, scopolamine, methscopolamine, oxyphenonium, papaverine; prostaglandins such as PGE₁, PGE₂, PGF₁ alpha., PGF₂ alpha. and PGA; antimicrobials

- 13 -

and antiparasitic agents such as penicillin, tetracycline, oxytetracycline, chloro-
tetracycline, chloramphenicol, thiabendazole, ivermectin, and sulfonamides; antimalarials
such as 4-aminoquinolines, 8-aminoquinolines and pyrimethamine; hormonal agents such
as dexamethasone prednisolone, cortisone, cortisol and triamcinolone; androgenic
5 steroids such as methyltestosterone, and fluoxmesterone; estrogenic steroids such as
17.beta.-estradiol, .alpha.-estradiol, estriol, .alpha.-estradiol 3-benzoate, and 17-ethynyl
estradiol-3-methyl ether; progestational steroids such as progesterone, 19-nor-pregn-4-
ene-3,20-dione, 17-hydroxy-19-nor-17-.alpha.-pregn-5(10)-ene-20-yn-3-one, 17.alpha.-
ethynyl17-hydroxy-5(10)-estren-3-one, and 9.beta., 10.alpha.-pregna-4,6-diene-3,20-
10 dione; sympathomimetic drugs such as epinephrine, phenylpropoudamine hydrochloride,
amphetamine, ephedrine and norepinephrine; hypotensive drugs such as hydralazine;
cardiovascular drugs such as procainamide, procainamide hydrochloride, amyl nitrite,
nitroglycerin, dipyredamole, sodium nitrate and mannitol nitrate; diuretics such as
chlorathiozide, acetazolamide, methazolamide, hydrochlorothiazide, amiloride
15 hydrochloride and flumethiazide, ethacrynic acid, furosemide; antiparasitics such as
bephenium, hydroxynaphthoate, dichlorophen and dapsone; and neoplastics such as
mechlorethamine, uracil mustard, 5-fluorouracil, 6-thioguanine and procarbazine; .beta.-
blockers such as pindolol, propranolol, practolol, metoprolol, oxprenolol, timolol, timolol
maleate, atenolol, alprenolol, and acebutolol; hypoglycemic drugs such as insulin,
20 isophane insulin, protamine zinc insulin suspension, globin zinc insulin, extended insulin
zinc suspension toblutamide, acetoexamide, tolazamide and chlorpropamide; antiulcer
drugs such as cimetidine; nutritional agents such as ascorbic acid, niacin, nicotinamide,
folic acid, choline, biotin, pantothenic acid, and vitamin B.sub.12 ; essential amino acids;
essential fats; eye drugs such as timolol, timolomaleate, pilocarpine, pilocarpine salts
25 such as pilocarpine nitriate, pilocarpine hydrochloride, dichlorphenamide, atropine,
atropine sulfate, scopolamine and eserine salicylate; histamine receptor antagonists such
as cimetidine; and electrolytes such as calcium gluconate, calcium lactate, potassium
chloride, potassium sulfate, sodium chloride, potassium fluoride, sodium fluoride, ferrous
lactate, ferrous gluconate, ferrous sulfate, ferrous fumurate and sodium lactate; and drugs
30 that act on .alpha.-adrenergic receptors such as clonidine hydrochloride.

- 14 -

Additional agents of interest include quinoline and naphthyridine carboxylic acids and related compounds, such as 1-ethyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid; 1-ethyl-1,4-dihydro-7-methyl-4-oxo-1,8-naphthyridine-3-carboxylic acid; 5-ethyl-5,8-dihydro-8-oxo-1,3-dioxolo[4,5-g]quinoline-7-carboxylic acid; 8-ethyl-5,8-dihydro-5-oxo-2-(1-piperazinyl)pyrido[2,3-d]pyrimidine-6-carboxylic acid; 9-fluoro-6,7-dihydro-5-methyl-1-oxo-1H,5H-benzo[ij]quinoxaline-2-carboxylic acid; 1-ethyl-1,4-dihydro-4-oxo-7-(4-pyridinyl)-3-quinolinecarboxylic acid; 1-ethyl-1,4-dihydro-4-oxo-[1,3]dioxolo[4,5-g]cinnoline-3-carboxylic acid; 9-fluoro-3-methyl-10-(4-methyl-1-piperazinyl)-7-oxo-2,3-dihydro-7H-pyrido[1,2,3-de][1,4]benzoxazine-6-carboxylic acid; 1-ethyl-6-fluoro-1,4-dihydro-7-(4-methyl-1-piperazinyl)-4-oxo-1,8-naphthyridine-3-carboxylic acid; 1-ethyl-6-fluoro-1,4-dihydro-7-(1-piperazinyl)-4-oxo-1,8-naphthyridine-3-carboxylic acid; 1-cyclopropane-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid; 1-methylamino-6-fluoro-1,4-dihydro-4-oxo-7-(4-methyl-1-piperazinyl)-3-quinolinecarboxylic acid; 1-(4-fluoro-1-phenyl)-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinolinecarboxylic acid; 1-(4-fluoro-1-phenyl)-6-fluoro-1,4-dihydro-4-oxo-7-(4-methyl-1-piperazinyl)-3-quinolinecarboxylic acid; 1-(4-fluoro-1-phenyl)-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-1,8-naphthyridine-3-carboxylic acid; and 1-ethyl-6-fluoro-1,4-dihydro-4-oxo-7-(3-ethylaminomethyl-1-pyrrolidinyl)-8-fluoro-3-quinolinecarboxylic acid.

20

Additional agents of interest include drugs which affect the respiratory tract such as, but not limited to, budesonide, enprofylline, tranilast, albuterol, theophylline, amoniphylline, brompheniramine, chlorpheniramine, promethazine, diphenhydramine, azatadine, cyproheptadine, terbutaline, metaproterenol, and isoproterenol; drugs which are antidepressants such as amiflamine, alaproclate, doxepin, trazedone, maprotiline, zimelidine, fluvoxamine; antipsychotic drugs such as haloperidol, thioridazine, trifluoperazine, MK-0212, and remoxipride; sedative hypnotic and antianxiety drugs such as triazolam, temazepam, chlorazepate, alprazolam, diazepam, fluorazepam, lorazepam, oxazepam, hydroxyzine, prazepam, meprobamate, butalbital, and chlorzoxazone; antiparkinson drugs such as benztropine and L-647,339; hormonal and steroidal drugs such as conjugated estrogens, diethylstilbestrol, hydroxy progesterone, medroxy progesterone, norethindrone, betamethasone, methylprednisolone, prednisone, thyroid

30

- 15 -

hormone, levothyroxine and MK-0621; antihypertensive and cardiovascular drugs such as isosorbide dinitrate, digoxin, nadolol, disopyramide, nifedipine, quinidine, lidocaine, diltiazam, verapamil, prazosin, captopril, enalapril, lisinopril, metyrosine, felodipine, tocanide, mexiletine, mecamlamine, and metyrosine; diuretic drugs such as
5 spironolactone, chlorthalidone, metolazone, triamterene, methyclothiazide, and indacrinone; antiinflammatory drugs such as ibuprofen, phenylbutazone, tolmetin, piroxicam, melclofenamate, auranofin, flurbiprofen and penicillamine; analgesic drugs such as acetaminophen, oxycodone, hydrocodone, and propoxyphene; antiinfective drugs such as cefoxitin, cefazolin, cefotaxime, cephalexin, nicarbazine, amprolium, ampicillin,
10 amoxicillin, cefaclor, erythromycin, nitrofurantoin, minocycline, doxycycline, cefadroxil, miconazole clotrimazole, phenazopyridine, clorsulon, fludalanine, pentizidone, cilastin, phosphonomycin, imipenem, arprinocid, and foscarnet; gastrointestinal drugs such as bethanechol, clidinium, dicyclomine, meclizine, prochlorperizine, trimethobenzamide, loperamide, ranitidine, diphenoxylate, famotidine, metoclopramide and omeprazole;
15 anticoagulant drugs such as warfarin, phenindione, and anisindione; and other drugs such as trientine, cambendazole, ronidazole, raxofenide, dactinomycin, asparaginase, nalorphine, rifamycin, carbamazepine, metaraminol bitartrate, allopurinol, probenecid, diethylpropion, dihydrogenated ergot alkaloids, nystatin, pentazocine, phenylpropanolamine, phenylephrine, pseudoephedrine, trimethoprim and mevinolin.

20

Therefore, the present invention provides for a delivery system comprising a hydrostatic couple in a solid matrix composition, the matrix composition containing one or more agents of interest with or without other pharmaceutical adjuvant(s). This delivery system ensures the release of one or more agents of interest in a controlled
25 manner, with a zero-order or near zero-order release kinetics, over a therapeutically practical time period. The delivery performance of the delivery system is only minimally affected by physiological variables in the gastrointestinal tract of human and animals.

By "hydrostatic couple" it is meant at least two components, for example, but not
30 limited to a group-A component (a hydrodynamic fluid-imbibing polymer) and a group-B component (a hydrostatic pressure modulating agent). Group-A component(s), are derived from at least one fluid-imbibing cross-linked polymer. Group-B components, are

- 16 -

derived from at least one rapid expansion source. Typically these components are placed within a matrix composition. When combined, and placed into a desired fluid environment, the two components act in consort to create a positive hydrostatic pressure that controls the release of one or more agents of interest from within the matrix composition to the adjacent environment. Preferably, the group-A component is dynamically permeable to the external fluids and solutes, and the group-B component is a rapidly expanding compound or provides a rapid expansion source upon interaction with the imbibed fluid.

Once liquid imbibition by capillary uptake is initiated it will continue until the entire solid polymer porous network of the matrix has been exhausted. During this process, the polymers as described herein, for example group-A polymers, undergo a dynamic non-limiting volume and mass increase due to a net influx of fluid. A desirable property of the group-A, hydrodynamic fluid-imbibing polymers, is their swelling ability characterized by weak intermolecular cohesive forces between the cross-linked chains when compared to the stronger intermolecular adhesive interaction between the fluid and polymer micropores. Liquid filled micropores are established during this process, thus allowing the diffusive efflux of solutes through these micropores. If enough solvating fluid is present, over time, the hydrodynamic fluid-imbibing polymers may start shedding.

The net influx of fluid in the polymer of this invention is dependent on both the pore size and microstructure of the polymer arrangement. Hydrodynamic polymers of this invention are cross-linked and the polymer molecular weight between crosslinks is indicative of their pore size. Differential hydrostatic pressure, and in consequence capillary uptake of fluid, within a polymer network that has a smaller pore size and or pronounced interpenetrating polymer structure will be greater in comparison to those polymer networks with a large pore size and fewer interpenetrating structures. In aqueous fluids these polymers will exhibit a linear or near-linear increase in volume and percent mass gain.

- 17 -

The fluid-imbibing polymers (group-A) of this invention are preferably cross-linked water insoluble polymers and can be in the state of dry powder, fine particle, granules or microcapsules.

5 As used herein, "hydrodynamic boundary" refers to any polymer or compound capable of relaxation upon exposure to the fluid media with a fixed and restricted radius of gyration due to, for example, extensive cross-linking. The molecular chains of these polymers are capable of forming insoluble boundaries or micropores containing water molecules sometimes referred in the art as microgels.

10

In the system of this invention the hydrodynamic boundary polymers are used in a concentration of about 4 to about 96 weight %, preferably about 60 to about 95, weight %, based on the total weight of the dosage unit.

15 Examples of suitable cross-linked group-A components (hydrodynamic fluid-imbibing polymer) capable of unlimited volume increase that can be used in this invention include but are not limited to:

- 20 • acrylic-acid polymers with cross-linking derived from allylsucrose or allylpentaerithritol, including water-insoluble acrylic polymer resins. Single compounds or a blend of compounds from this group of polymers include for example, but not limited to Carbopol.RTM.971-P, Carbopol.RTM.934-P, Carbopol.RTM.974P and Carbopol.RTM.EX-507 (GF Goodrich, or any other commercially available brand with similar properties, may be used). Preferably, 25 the acrylic-acid polymers have a viscosity from about 3,000 centipoise to about 45,000 centipoise at 0.5% w/w concentration in water at 25°C, and a primary particle size range from about 3.00 to about 10.00 microns in diameter, as determined by Coulter Counter;
- 30 • highly cross-linked or lightly cross-linked starch derivatives crosslinked by Epichlorhydrin or Phosphorous oxychloride (POCl_3) or Sodium trimetaphosphate are also suitable for use in the hydrostatic delivery system described herein either singly or in blends;

- 18 -

- polyglucans such as amylose, dextran, pullulan succinates and glutarates containing diester –crosslinks either singly or in blends;
- diether crosslinked polyglucans such as those disclosed in U.S. 3,208,994 and 3,042,667 (which are incorporated herein by reference);
- 5 • crosslinked polyacrylate resins such as, but not limited to, potassium polyacrylate; and
- water swellable crosslinked polymer compositions of crosslinked polyethylenimine and or crosslinked polyallylamine.

10 It is further contemplated that mixtures of the above compounds may be used as a group-A component.

Examples of methods of preparation, for example of Carbopol.RTM.934-P, - a polymer of acrylic acid lightly cross-linked with polyallyl ether of sucrose having an
15 average of 5.8 allyl groups per each sucrose molecule, has been disclosed in U.S. 2,909,462; 3,033,754; 3,330,729; 3,458,622; 3,459,850; and 4,248,857 (which are incorporated herein by reference). When Carbopol.RTM.971-P is used, the preferred viscosity of a 0.5% w/w aqueous solution is 2,000 centipoise to 10,000 centipoise. More preferably, the viscosity of a 0.5% w/w aqueous solution is 3,000 centipoise to 8,000
20 centipoise. When Carbopol.RTM.934-P is used, the preferred viscosity of a 0.5% w/w aqueous solution is 20,000 centipoise to 60,000 centipoise, more preferably, the viscosity of a 0.5% w/w aqueous solution is 30,000 centipoise to 45,000 centipoise

Cross-linked starch derivatives (crosslinked by Epichlorhydrin or Phosphorous
25 oxychloride (POCl.sub.3) or Sodium trimetaphosphate) include high amylose starch containing varying degrees of crosslinking. These compounds and their methods of preparation are known in the art, for example, U.S. 5,807,575 and U.S. 5,456,921 (which are incorporated herein by reference), and Rutenberg and Solarek (M.W. Rutenberg and D. Solarek, "Starch derivatives: production and uses" in Starch Chemistry and
30 Technology, 2nd Edition, Chapter X, Pages 311-379, R.L. Whistler, J.N. BeMiller and E.F. Paschall, Academic Press, 1984; which is incorporated herein by reference).

- 19 -

Water-insoluble polyglucans such as amylose, dextran, pullulan succinates and glutarates containing diester crosslinks also exhibit water imbibition properties suited for group-(A) components of this invention. The methods and manufacture of these compounds have been disclosed in U.S. 4,002,173 (which is incorporated herein by reference). These individual polyglucans or blends derived therefrom can be used to satisfy group-(A) component.

Crosslinked polyacrylate resins such as potassium polyacrylate having sufficiently low water content and capable of being powdered can be used in this invention (see U.S. 4,654,393 and 4,954,562, which are incorporated herein by reference). These polyacrylate resins are highly water absorbing and insoluble.

As used herein, "hydrostatic pressure-modulating agent" refers to one or more compounds that expand rapidly upon exposure to fluid, for example, a rapid expansion polymer (a group-B component), serve as an expansion source (see below), or a combination of a rapid expansion polymer and an expansion source.

Hydrophilic cross-linked polymers, known in the art, as super disintegrants can exhibit rapid expansion upon exposure to aqueous media. These polymers are capable of rapid capillary uptake of water and a limiting volume expansion. The limited volume expansion is characterized by an intermolecular cohesive force between the polymer molecules that is stronger in comparison to weaker intermolecular adhesive forces, between the solvent molecules and the polymer pore structure. The rate and extent of volume expansion is dependent on the particle size and nature of the polymer cross-links. A polymer with small particle sizes contains fewer cross-links per unit particle and expands faster, but to a lesser extent, in comparison to the same polymer with larger particle sizes. The rate and extent of expansion in these polymers is much faster than their inherent rate and extent of water imbibition.

Examples of rapid expansion polymers suitable as group-B components of this invention include, but are not limited to:

- 20 -

- single compounds or combinations derived from cross-linked N-vinyl-2-pyrrolidone (PVP) selected from a group of chemically identical polyvinylpolypyrrolidone such as Polyplasdone.RTM.XL, Polyplasdone.RTM.XL-10, Polyplasdone.RTM.INF-10 (International Specialty Products). Preferably, the cross-linked N-vinyl-2-pyrrolidone has a particle size from about 9 microns to about 150 microns; and
- cross-linked cellulose derivatives selected from a group of hydrophilic compounds such as cross-linked carboxymethyl cellulose (for example croscarmellose), sodium starch glycolate or a combination thereof.

Therefore, the present invention provides a hydrostatic delivery system comprising a hydrostatic couple, wherein the hydrostatic couple comprises at least one hydrodynamic fluid-imbibing polymer, and at least one hydrostatic pressure modulating agent. Preferably, the hydrodynamic fluid-imbibing polymer is a cross-linked polymer with a swelling capacity in a fluid environment of between about 1 weight% to about 3000 weight%. By swelling capacity it is meant the percentage gain in mass as result of water imbibition, for example as determined using:

$$\text{(Mass at time (t) - Initial Dry Mass) / Initial Mass .}$$

Preferably, the hydrodynamic fluid-imbibing, cross-linked polymer is present from about 4 weight% to about 96 weight% of the total formulation. Furthermore, the hydrostatic pressure modulating agent is preferably a cross-linked, rapidly swelling polymer with a swelling capacity in fluid environment of between about 0.5 weight% to about 500 weight%. Preferably, the cross-linked, rapidly swelling polymer (the hydrostatic pressure modulating agent) is from about 0.5 weight% to about 50 weight% of the total formulation. However, hydrostatic delivery systems comprising greater than 50 weight %, for example, up to 80 weight%, are also contemplated, depending upon the rate of delivery required, and the drug being delivered.

- 21 -

The present invention also pertains to a hydrostatic delivery system comprising a hydrodynamic polymer and a hydrostatic pressure modulating agent, wherein the hydrodynamic polymer and the hydrostatic pressure modulating agent are present at a ratio from about 99:1 to about 1:99. Preferably, the hydrodynamic polymer and the hydrostatic pressure modulating agent are present at a ratio from about 99:1 to about 50:50.

An "expansion source" is a hydrostatic pressure-modulating agent that is able to create a hydrostatic pressure within the hydrostatic delivery system of the present invention. For example, which is not to be considered limiting, an expansion source may be an alkaline agent capable of releasing a gas, or causing a fluid to effervesce, when exposed to a proton source such as an acidic agent or water. In this manner, the alkaline agent can serve as an expansion source capable of creating a hydrostatic pressure within the hydrostatic delivery system of the present invention. The alkaline agent can be carbon dioxide gas precursor, an oxygen gas precursor or a chlorine dioxide gas precursor.

Alkaline agents of this invention can be selected from, but are not limited to, carbon dioxide precursors such as carbonates, sesquicarbonates and hydrogencarbonate salts of potassium, lithium, calcium, sodium, ammonium, L-lysine carbonate, arginine carbonate, sodium glycine carbonate and sodium amino acid carbonate. The alkaline agents can also be obtained from a group of oxygen gas precursor such as, but not limited to, anhydrous sodium perborate, effervescent perborate, sodium perborate monohydrate, sodium percarbonate and sodium dichloroisocyanurate. Chlorine dioxide (ClO_2) precursor compounds such as sodium hypochlorite can also be used as alkaline agents in applications such as cleansing operations.

Therefore, the present invention also provides for a hydrostatic delivery system wherein the hydrostatic pressure modulating agent comprises an expansion source, selected from the group consisting of a carbon-dioxide precursor, an oxygen precursor, and a chlorine dioxide precursor. Preferably, when the hydrodynamic polymer comprises a carbon dioxide precursor, oxygen precursor or chlorine dioxide precursor, the hydrodynamic polymer and the hydrostatic pressure modulating agent are present in a

- 22 -

ratio from about 99:1 to about 50:50 by weight, preferably, from about 99:1 to about 70:30 by weight.

As used herein, the term "acidic agent" refers to any compound or material that can serve as a proton source and can react with the alkaline agent of the invention to form a gas. The acidic agent can have more than one acid functional group, that is, more than one dissociation constant. The acidic agent can be any organic or inorganic acid in the free acid, acid anhydride and acid salt form. Preferably, the acidic agent is in a solid state at ambient temperatures, is not harmful to animals including man, and exhibits a pH of about 4.6 or lower when saturated into water at room temperature. Also included as acidic agents are acid alkali metal salts (e.g. sodium salt, potassium salt, etc.).

Examples of an acidic agent include, but are not limited to, citric acid, tartaric acid, fumaric acid, maleic acid, malic acid, lactic acid, succinic acid, adipic acid, glycolic acid, alpha hydroxy acids, ascorbic acid, amino acids and their alkali hydrogen acid salts, as well as alkali acid metal salts of acid substances such as phosphoric acid and pyrophosphoric acid or other inorganic acids provided those salts are solid at room temperature. The preferred type of acidic agent possess a relatively large acid dissociation constant (10^3 or more) and a low hygroscopicity (critical humidity at 30°C. is 40% RH or more).

The ratio of the acidic agent and alkaline agent can be determined according to the amount of gas required to effect a desirable hydrostatic pressure. When the two compounds, i.e. the acidic and alkaline agents, are mutually reactive, it is preferable, although not necessary, that they react completely. Therefore, a ratio of components that provides for equal amounts of reaction equivalents is preferred. For example, if the acid used is diprotic, then either twice the amount of a mono-reactive carbonate alkaline agent, or an equal amount of all-reactive alkaline agent should be used for complete neutralization. The amount of alkaline agent can be increased if it is desired to increase the hydrostatic pressure of the delivery system. If the alkaline agent is a carbon dioxide precursor, the amount of the precursor within the delivery system varies from about 0.5 to about 70 weight%, preferably 2 to about 30 weight%, of the formulation. Preferably

- 23 -

the acidic agent and a alkaline agent, for example a carbon dioxide precursor, are solid, for example in a powdery or granular state.

5 The hydrostatic couple as described herein, comprises a mixture of at least one group-A, hydrodynamic boundary compound, and at least one group-B, hydrostatic pressure-modulating agent. The combination thus formed is capable of establishing a positive hydrostatic differential pressure against the fluid environment.

10 The ratio of the group-A and group-B compounds in the hydrostatic couple is determined according to the amount of hydrostatic pressure required to produce a desired volume efflux. This ratio is also related to the hydrostatic pressure required to reach equilibrium "steady state" volume for the delivery system. The amount of hydrostatic couple in a particular delivery system will depend on the saturation solubility of the agent of interest to be delivered, the desired rate and the duration of such release from the
15 delivery system.

 The rate of release of the agent of interest may be selected considering several variables, for example, but not limited to the solubility of the agent of interest, and pharmacological activity of the agent of interest. For example, with decreased solubility
20 or a low pharmacological activity, of an agent of interest, a faster release of the agent from the hydrostatic couple may be desired. Likewise, with a soluble agent of interest, or an agent that exhibits a high degree of pharmacological activity, a slower release of the agent from the delivery system may be desired, for example, but not limited to, using a hydrostatic couple as provided in Formula 2. It is to be understood, however, that the
25 formulation of the hydrostatic couple may be varied as required to obtain a desired rate of release of an agent of interest.

 The present invention also provides for a delivery system that utilizes capillarity as a means of fluid imbibition, and differential volume expansion of the hydrostatic
30 couple to create a non-osmotic hydrostatic pressure. The resultant hydrostatic pressure produces the driving force for the controlled release of the agent of interest. The hydrodynamic boundary polymers of the hydrostatic couple, for example the group-A

- 24 -

component(s), create a hydrodynamic boundary which is capable of moving, and is in dynamic equilibrium with, the components of the hydrostatic couple.

5 A further aspect of the present invention provides for a solid pharmaceutical dosage form, for example, but not limited to a matrix compact suitable for oral administration wherein the delayed release is brought about by use of suitable excipients that are industrially available, non-toxic and easy to process. The pharmaceutical dosage form includes, for example, but not limited to, compressed tablets, granules, pellets, suspensions, extrusion spheroids or compacts obtained by direct compression, wet
10 granulation, dry granulation, hot melt granulation, microencapsulation, spray drying, and extrusion methods as would be evident to one of skill in the art. Other solid dosage forms such as hard gelatin capsules can also be derived from dry blends, granulations, suspensions, spheroids, pellets, tablets and combinations therefrom, as are commonly known in the art.

15

The pharmaceutical dosage form may also include excipients as required, for example, but not limited to, viscosity enhancer(s), enteric polymer(s), pH-specific barrier polymer(s), diluent(s), anti-adherent(s), glidant(s), binder(s), solubilizer(s), channeling agent(s), wetting agent(s), buffering agent(s), flavorants, adsorbents, sweetening agent(s),
20 colorant(s) and lubricant(s) :

25

The dosage forms and delivery system taught herein may be used in pharmaceutical, veterinary, food, pesticidal, horticultural, herbicidal, agricultural, cosmetic, industrial, cleansing, and confectionery applications.

30

Formulations incorporating the solid dosage forms can further include one or more additional adjuvants, which can be chosen from those known in the art including flavors, colors, diluents, binders, plasticizers, fillers, surfactant, solubilizers, stabilizers, compaction enhancers, channeling agents, glidants, lubricants, coating polymers and anti-adherents.

Hydrostatic Delivery System

- 25 -

When the hydrostatic delivery system of the present invention comes into contact with an external fluid of the environment, such as water or other biological fluid, the water of fluid is imbibed into the core of the delivery system in part by capillary hydration due to the hydrostatic couple. The volume of the hydrodynamic boundary, group-A components of the system increases due to a net inflow into the polymer structure. Concurrently, the rapidly expanding hydrostatic-pressure modulating agent (group-B component) also increases in volume. However due to the differential rates and extents of volume increase of the individual polymers, a positive differential hydrostatic pressure builds up within the delivery system. An expansion source also exerts pressure against the polymeric micropores of the cross-linked group-A polymer and thus produces a net differential pressure. The differential fluid pressure therefore is in part derived from the hydrostatic couple typically arising from the net effect of two dynamically independent processes contributed by each component of the couple.

Without wishing to be bound by theory, at a given point of net imbibed water, there is a given ratio of the number of hydrated group-A particles and expanded particles or molecules of the rapid expansion-group-B particles, that creates a positive differential pressure. This hydrostatic pressure acts against the influx of water and at some point the inflow of water will equal the outflow of water. The resultant hydrostatic volume efflux overwhelms the passive diffusive volume efflux within the delivery system. When the inflow and outflow of water become equal, the system manifests a dynamic constant volume and surface area. This results in a steady state release of solved or partially solved particles of the agent of interest along with any other adjuvant.

Without wishing to be bound by theory, the kinetics of volume fluxes due to the hydrostatic couple may be explained as follows: Upon contact with an external fluid, due to surface tension of the liquid and surface free energy at the solid-liquid interface, the liquid wets the pores of the hydrostatic couple components. Fluid enters into the porous structure of the components due to hydrostatic capillary action and surface tension. This imbibition continues so long as there is a pressure difference as a result of density differences between the imbibed liquid and the vapor (or air) within the solid polymer

- 26 -

network of the hydrostatic couple. Therefore water inflow will continue until all solid components and polymer pores have been hydrated.

The volume influx from capillary action (k), leading to volume increase is
 5 $(dV/dt)_k$. This rise in volume is opposed by the diffusive flux, $(dM/dt)_d$, due to chemical potential gradient of dissolved solutes or agent of interest, and is driven by passive diffusion. The diffusive flux $(dM/dt)_d$ is equal to the product of the passive volume efflux and the concentration (C) of the solute; $(dV/dt)C_e$. In the absence of the hydrostatic pressure modulating agent(s), the net volume flux $(dV/dt)_T$ is given by the following
 10 equation (equation 1):

$$\text{Equation 1: } dV/dt_T = dv/dt_K - [dV/dt \times C]_e$$

The rate of volume influx is substantially greater than the diffusive rate of efflux, thus a
 15 constant volume increase is observed (see Figure1).

In the hydrostatic delivery system of this invention, the presence of a hydrostatic couple creates a positive hydrostatic pressure within the delivery system as a result of the differential rates of volume expansion between the group-A and group-B components.
 20 This differential pressure opposes the volume influx of the imbibed fluid and reduces the volume gain of delivery system. The volume efflux due the hydrostatic pressure $(dV/dt)_h$, is substantially greater than the contribution to volume efflux as a result of passive diffusive flux. At an optimal level, which may be determined by a mathematically predictable ratio of the components of the hydrostatic couple, the rate of volume efflux
 25 approaches and eventually equals the rate of volume influx. This represents a dynamic steady state with a zero net increase in volume and a constant surface area of the delivery system (see Figure2). The one or more agents of interest that are dissolved or partially solved within the delivery system, are thus released at a rate determined by the total (net) efflux controlled and determined by hydrostatic pressure within the delivery system. The
 30 insignificance of the passive diffusive contribution to the net volume efflux proffers a delivery system whose performance is independent of the chemical concentration

- 27 -

gradient of the agent of interest. If the volume within the delivery system is such that the total concentration of the agent of interest is above its saturation concentration, the resultant release of the agent of interest will exhibit a zero or near zero order kinetics. The net volume flux in the hydrostatic delivery system is represented by the following equation (equation 2):

Equation 2:
$$dV/dt_T = dV/dt_K - [dV/dt_h + dV/dt_d \times C]_e$$

$$dV/dt_T = dV/dt_K - [dV/dt_h \times C]_e$$

The dynamic fluid profile of a delivery system of the present invention comprising a hydrostatic couple as described herein is presented in Figure 2. Following an initial increase in the dynamic volume of the tablet, the volume remains stable over time wherein the influx of fluid is equal to the efflux of caffeine. This represents a controlled increase in the dynamic profile of a tablet, which reaches and maintains a maximum volume after a period of time (depending upon the ratio of the hydrodynamic fluid-imbibing polymer, to hydrostatic pressure modulating agent). Figures 4 - 6 show a corresponding drug release (dissolution profile) for a range of formulations comprising a hydrostatic couple of the present invention and a range of concentrations of active ingredients. The dissolution profiles shown in Figure 4-6 each display a linear, zero-order release of an agent of interest for over 16 hours.

The above description is not intended to limit the claimed invention in any manner, furthermore, the discussed combination of features might not be absolutely necessary for the inventive solution.

The present invention will be further illustrated in the following examples. However it is to be understood that these examples are for illustrative purposes only, and should not be used to limit the scope of the present invention in any manner.

Examples

- 28 -

Example 1: Preparation of hydrostatic couple

The hydrostatic couple is prepared either by dry screening the components of group- A and group-B compounds followed by geometric blending of components to form an intimate mixture, or granulating the components to form discrete aggregates. The agent of interest is prepared and combined with the hydrostatic couple. This may be achieved by blending or further re-granulation to homogeneity. The resulting mixture is compressed on a B or D type tablet press to form a compact. Such compacts may be further coated if desired using standard techniques as known within the art.

A manufacturing process for a hydrostatic delivery system in form of a matrix tablet involves the following general steps:

- a) Preparation of group-A component(s);
- b) Preparation of group-B component(s);
- c) Preparation of agent of interest formulation;
- d) Blending a)-c);
- e) Compressing the blend into a suitable compact;
- f) Coating the compact with a suitable polymer.

Preparation of group- A granules

Granulate the selected item(s) of group-A component with 100% Isopropyl alcohol or suitable granulating fluid in a high-shear mixer-granulator. Wet screen in an Oscillating Granulator or suitable granulator equipped with mesh # 16 or # 20 or other suitable size. Dry the wet screened moist granules in a convection oven set at room temperature for 60-90 minutes. Dry screen in a Comill or suitable dry granulator (mesh 0.5-mm, 200 –250 rpm). Dry further at 30-35°C for 3 – 4 hours. Sieve in a Rotap for 3-4 minutes (screen # 60/100).

Preparation of group-B granules

- 29 -

Granulate the selected item(s) of group-B component with 100% Ethanol or other suitable solvent in a high-shear mixer-granulator. Wet screen in an Oscillating Granulator or suitable granulator equipped with mesh # 16 or # 20 or other suitable size. Dry the wet screened moist granules in a convection oven set at 30-35°C for 90 – 120 minutes. Sieve
5 in a Rotap for 3-4 minutes (screen # 16/60).

Preparation of active component blend comprising an agent of interest

The active compound blend may be prepared using any suitable method as would
10 be known to one of skill in the art for example those disclosed in Lachman et al. (The Theory and Practice of Industrial Pharmacy, by L. Lachman, H. Lieberman and J. Kang. 3rd Edition, Lea & Febiger 1986). This may involve combination of the agent of interest with an excipient, or granulation, or microencapsulation or other suitable method.

15 Blending & Compressing

Group-A and group-B granules are uniformly blended by geometric dilution in a Paterson-Kelly (PK) twin shell blender. After discharge, the active component blend is serially mixed with the blend composition of group-A and group-B components for
20 about 6-8 minutes. The final composition is discharged and compressed into a suitable sized compact.

Coating

25 If desired, the compressed compact may be coated with an aqueous or solvent based polymer solution. The choice of polymer, solvent and plasticizer may vary as required and it is dependent on the desired outcome. The polymer may be a functional coating such as a pH-dependent enteric polymer or a non-functional coating such as a hydrosoluble polymer for esthetics.

30

Example 2:

- 30 -

Two drug delivery systems were prepared in order to compare the kinetics of drug release and changes in dynamic volume profile of the delivery system (tablet). The control comprised a prior art formulation, (Prior Art, Table 1) and the second delivery system comprised the hydrostatic couple (components listed in Table 1) as prepared using the method of Example 1. The agent of interest in both delivery systems was caffeine.

Table 1: Components of Prior art and hydrostatic couple formulations

Component	Component	Amount per tablet (mg)
Prior art		
Active agent	Caffeine Anhydrous USP	160
Control Release polymer	Carbopol 971P USP/NF	224
Flow Promoter	Colloidal Silicon Dioxide	12
Lubricant	Magnesium Stearate	4
Hydrostatic couple		
Active agent	Caffeine Anhydrous USP	70
Hydrostatic couple		
Group-A	Carbopol 971P	280
Group-B	Crospovidone XL-10	8
Flow promoter	Colloidal Silicon Dioxide	4.3
Lubricant	Magnesium Sterate	3.67

The prepared delivery systems were placed within PBS at pH7.0 in a Type II USP 24 Dissolution apparatus at 37°C (±0.5) using a paddle speed of 50 rpm. Caffeine release from the delivery systems were measured over time. Caffeine release was determined spectrophotometrically @ 272nm.

Method for Measuring Dynamic Volume Change due to Fluid Imbibition

- 31 -

The dynamic volume change of a fluid-imbibing or swelling tablet was measured by computation of the density of the swollen tablet and its mass. The basic relationship is:

$$V_t = M_t / D$$

5

Where V_t is the volume at a given time; M_t is the mass of swollen tablet at a given time; and D , is the density of the swollen tablet

To obtain dynamic volume values, the same tablet undergoing swelling in the fluid media was removed from the dissolution media at regular (pre-fixed) time intervals, weighed in air (to obtain its mass) and weighed submerged in the fluid media (to obtain its buoyancy). The tablet is immediately returned to the dissolution media where swelling resumes. The time lapse between removal from the fluid media and its return to the media is kept constant and short in order to minimize errors due to excessive dehydration. This time interval is typically not more than 30 seconds.

The density of the swollen tablet is obtained by calculating:

$$\rho_2 = (A/P) * \rho_0$$

20

where, ρ_2 is the density of the swollen tablet; ρ_0 is the density of the fluid media.

Equipment & Materials for Dynamic Volume Measurement

Swelling & Drug Dissolution Measurements

25

USP Dissolution Apparatus Type II (Paddle)

Settings: Rotational Speed: 40 – 50 rpm

Temperature: 37° C +/- 0.5° C

PBS buffer pH 7.00 (or suitable buffer at a desired pH).

30

Dynamic Volume Measurements

Mettler-Toledo Density Determination Kit (for liquids and Solids) Model # 33360

Media: PBS buffer pH 7.00 or suitable buffer at a desired pH.

- 32 -

Figure 1 shows a plot of the dynamic volume profile of a prior art formulation, demonstrating a linear volume increase associated with a hydrodynamic polymer (Group-A component). The corresponding drug release (dissolution profile) for this formulation is shown in Fig 3. A rapid release (exponential) of an agent of interest from the prior art delivery system, reaching a maximum release rate after about 3.5 to 4 hours is evident in Figure 3. This is the typical Fickian release manifested by prior art compositions using group A- type components as the control release polymer. With this delivery system, the rate of efflux of an agent of interest is due to passive diffusion and is substantially less than the rate of influx of the fluid. Consequently, the rate of release of an agent of interest is dependant on the chemical potential and concentration of the agent.

The dynamic fluid profile of a delivery system of the present invention comprising a hydrostatic couple provided in Table 1 is presented in Figure 2. Following an initial increase in the dynamic volume of the tablet, the volume remains stable over time wherein the influx of fluid is equal to the efflux of caffeine. This represents a controlled increase in the dynamic profile of a tablet, which reaches and maintains a maximum volume after a period of time (depending upon the ratio of the hydrodynamic fluid-imbibing polymer, to hydrostatic pressure modulating agent). Figure 4 shows the corresponding drug release (dissolution profile) for a formulation comprising a hydrostatic couple of the present invention, and displays a linear, zero-order release of an agent of interest for over 16 hours. Figures 2 and 4 demonstrate how the volume increase in a delivery system comprising a hydrostatic couple is reduced, resulting in an increased and continuous efflux rate. Because the rate of efflux of the agent of interest is independent on the concentration of the agent but dependent on the hydrostatic pressure within the delivery system, the kinetics of agent release is zero-order.

Examples 3-6:

In these examples hydrostatic delivery systems for extended release formulation of various therapeutic agents are presented. Two formulations (Formula 1 and Formula 2) are used to illustrate how the hydrostatic couple as described herein can be used to

- 33 -

achieve zero-order kinetics and predictably different release rates. Formula 1 exhibits faster rates of drug release than that observed with formula 2. The rate of release of the agent of interest may be selected considering several variables, for example, but not limited to the solubility of the agent of interest, and pharmacological activity of the agent of interest. For example, with decreased solubility of an agent of interest, faster release of the agent may be desired, such as that provided by, but not limited to, Formula 1. In the case of a soluble agent of interest, slower release of the agent from the delivery system may be desired, for example, but not limited to, using a hydrostatic couple as provided in Formula 2. It is to be understood, however, that the formulation of the hydrostatic couple may be varied as required to obtain a desired rate of release of an agent of interest.

Example 3 : Extended Release Theophylline 80 mg**Table 2: Extended Release Theophylline**

Components	Formula-1	Formula-2
Theophylline USP	80.00 mg	80.00 mg
Carbopol 971P NF	320.00 mg	320.00 mg
Crospovidone XL-10	6.40 mg	0.00 mg
Crospovidone NF-10	0.00 mg	6.40 mg
Sodium Lauryl Sulphate NF	4.00 mg	4.00 mg
Colloidal Silicon Dioxide NF	3.00 mg	3.00 mg

Example 4: Extended Release Nifedipine 60 mg**Table 3: Extended Release Nifedipine**

Components	Formula-1	Formula -2
Nifedipine USP	60.00 mg	60.00 mg
Carbopol 971P NF	171.00 mg	171.00 mg
Carbopol 934P NF	9.00 mg	9.00 mg
Crospovidone XL-10	3.60 mg	0.00 mg

- 34 -

Crospovidone INF-10	0.00 mg	3.60 mg
Cyclodextrin NF	2.00 mg	2.00 mg
Sodium Lauryl Sulphate	1.50 mg	1.50 mg
Colloidal Silicon Dioxide NF	3.00 mg	3.00 mg

5

Example 5 : Extended Release Diltiazem 60 mg**Table 4: Extended Release Diltiazem**

10	Components	Formula-1	Formula-2
	Diltiazem USP	60.00 mg	60.00 mg
	Carbopol 971P NF	171.00 mg	171.00 mg
	Crospovidone XL-10	3.60 mg	0.00 mg
	Crospovidone INF-10	0.00 mg	3.60 mg
15	Cyclodextrin NF	2.00 mg	2.00 mg
	Colloidal Silicon Dioxide NF	3.00 mg	3.00 mg

Example 6: Extended Release Buspirone hydrochloride 20 mg

20

Table 5: Extended Release Buspirone hydrochloride

25	Components	Formula-1	Formula-2
	Buspirone Hydrochloride USP	20.00 mg	20.00 mg
	Carbopol 971P NF	171.00 mg	171.00 mg
	Crospovidone XL-10	3.60 mg	0.00 mg
	Crospovidone INF-10	0.00 mg	3.60 mg
	Cyclodextrin NF	2.00 mg	2.00 mg
	Colloidal Silicon Dioxide NF	3.00 mg	3.00 mg

30

Example 7: Extended Release Rantidine Hydrochloride 60mg

- 35 -

The preparation of an extended release formulation composition comprising Rantidine Hydrochloride (Table 6) was prepared as described herein (Example 1). The release of Rantidine hydrochloride was measured in a USP dissolution apparatus type II dissolution apparatus under the following conditions:

- 5 Paddle speed: 50 rpm
 Temperature: 37°C (±0.5)
 Media: PBS buffer pH 7.0
 Volume: 900 ml
 Sampling Duration: 24 hours.

- 10 Ranitidine hydrochloride release was determined spectrophotometrically @ 322nm. The results for this experiment are shown in Figure 5.

Table 6: Composition of Rantidine Hydrochloride Tablets

Component		Amount
15 Model drug	Ranitidine Hydrochloride	60.00mg
Hydrostatic Couple		
Group-A	Carbopol 971P	203.66mg
Group-B	Crospovidone XL-10	1.54mg
Other excipients		
20 Binder (pH sensitive)	Hydroxyl Propoyl Methyl Cellulose Phthalate	4.00mg
Lubricant	Magnesium stearate	4.00mg

- 25 As shown in Figure 5, the dissolution profile for Ranitidine Hydrochloride from a formulation comprising a hydrostatic couple of the present invention, displays a linear, zero-order release of an agent of interest for over 16 hours.

Example 8: Extended Release Tramadol Hydrochloride 200mg

- 30 Extended release formulations comprising Tramadol hydrochloride (200 mg; composition outlined in Table 7) were prepared as described in Example 1. Extended release of Tramadol hydrochloride was measured in a USP dissolution apparatus type II dissolution apparatus under the following conditions:

- 36 -

Paddle speed: 50 rpm

Temperature: 37°C (±0.5)

Media: PBS buffer pH 7.0

Volume: 900 ml

5 Sampling Duration: 24 hours

Tramadol hydrochloride release was determined spectrophotometrically @ 275nm. The results of this experiment are shown in Figure 6.

10 **Table 7: Composition of Tramadol Hydrochloride Tablets**

Component		Amount
Model drug	Tramadol Hydrochloride	200.00mg
Hydrostatic Couple		
Group-A	Carbopol 971P	200.50mg
15 Group-B	Crospovidone XL-10	1.20mg
Other excipients		
Lubricant	Magnesium stearate	2.00mg

20 The results shown in Figure 6, demonstrate that the dissolution profile for Tramadol hydrochloride from a formulation comprising a hydrostatic couple of the present invention, displays a linear, zero-order release of an agent of interest for over 16 hours.

25 All citations are herein incorporated by reference.

The present invention has been described with regard to preferred embodiments. However, it will be obvious to persons skilled in the art that a number of variations and modifications can be made without departing from the scope of the invention as
30 described herein.

- 37 -

THE EMBODIMENTS OF THE INVENTION IN WHICH AN EXCLUSIVE PROPERTY OF PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A hydrostatic delivery system comprising a hydrostatic couple and an agent of interest.
2. The hydrostatic delivery system of claim 1 wherein said hydrostatic couple comprises, at least one hydrodynamic fluid-imbibing polymer, and at least one hydrostatic pressure modulating agent.
3. The hydrostatic delivery system of claim 2, wherein said hydrodynamic fluid-imbibing polymer is a cross-linked polymer having a swell capacity in a fluid environment of between about 1 weight% to about 3000 weight%.
4. The hydrostatic delivery system according to claim 3, wherein said cross-linked polymer is present from about 4 weight% to about 96 weight% of the total formulation.
5. The hydrostatic delivery system according to claim 2, wherein said hydrostatic pressure modulating agent is a cross-linked, rapidly swelling polymer having a swell capacity in a fluid environment of between about 0.5 weight% to about 500 weight%.
6. The hydrostatic delivery system according to claim 5, wherein said cross-linked polymer present from about 0.5 weight% to about 50 weight% of the total formulation.
7. The hydrostatic delivery system according to claim 2, wherein said hydrostatic pressure modulating agent, further comprises an expansion source.
8. The hydrostatic delivery system according to claim 7, wherein said expansion source is selected from the group consisting of a carbon-dioxide precursor, an oxygen precursor, and a chlorine dioxide precursor.

- 38 -

9. The hydrostatic delivery system according to claim 3, wherein said hydrodynamic polymer and said hydrostatic pressure modulating agent are present at a ratio from about 99:1 to about 50:50 by weight.

10. The hydrostatic delivery system according to claim 3, wherein said hydrodynamic polymer is a carbon dioxide precursor, an oxygen precursor or a chlorine dioxide precursor, said hydrodynamic polymer and said hydrostatic pressure modulating agent present in a ratio from about 99:1 to about 70:30 by weight.

11. The hydrostatic delivery system according to claim 1 wherein the agent of interest comprises a plurality of discrete active particulates.

12. The hydrostatic delivery system according to claim 2, wherein said hydrodynamic fluid-imbibing polymer comprises one or more of the compounds selected from the group consisting of:

- i) an acrylic-acid polymer cross-linked with allylsucrose or allylpentaerythritol;
- ii) one or more starch derivatives cross-linked by Epichlorhydrin, Phosphorous oxychloride (POCl_3), or Sodium trimetaphosphate;
- iii) a polyglucan;
- iv) a crosslinked polyacrylate resin;
- v) a crosslinked polyethylenimine;
- vi) a crosslinked polyallylamine, and combinations thereof.

13. The hydrostatic delivery system according to claim 2, wherein the hydrostatic pressure modulating agent comprises one or more of the compounds selected from the group consisting of:

- i) homopolymers of cross-linked N-vinyl-2-pyrrolidone;
- ii) a rapidly expanding cross-linked cellulose derivative; and combinations thereof

- 39 -

14. The hydrostatic delivery system according to claim 12, wherein said acrylic-acid polymer is selected from a group consisting of Carbopol.RTM.971P, Carbopol.RTM.934P, Carbopol.RTM.971P, Carbopol.RTM. EX507, and a combination thereof.

15. The hydrostatic delivery system according to claim 14, wherein said acrylic-acid polymer has a viscosity from about 3,000 centipoise to about 45,000 centipoise at 0.5% w/w concentration in water at 25°C.

16. The hydrostatic delivery system according to claim 15, wherein said acrylic-acid polymer has a primary particle size range from about 3.00 to about 10.00 microns in diameter.

17. The hydrostatic delivery system according to claim 12, wherein said polyglycan is selected from the group consisting of amylose, dextran, pullman succinate containing diester or diether crosslinks, pullman glutarates containing diester or diether crosslinks, and a combination thereof.

18. The hydrostatic delivery system according to claim 13, wherein said homopolymers of cross-linked N-vinyl-2-pyrrolidone are selected from the group consisting of Polyplasdone.RTM.XL, Polyplasdone.RTM. XL-10, Polyplasdone.RTM. INF-10, and a combination thereof.

19. The hydrostatic delivery system according to claim 16, wherein said cross-linked N-vinyl-2-pyrrolidone has a particle size from about 9 microns to about 150 microns.

20. The hydrostatic delivery system according to claim 13, wherein said rapidly swelling cross-linked cellulose derivative is selected from the group consisting of cross-linked carboxymethyl cellulose, sodium starch glycolate, and a combination thereof.

21. The hydrostatic delivery system according to claim 8, wherein said carbon dioxide precursor is selected from the group consisting of carbonates, sesquicarbonate,

- 40 -

hydrogencarbonate, potassium carbonate, lithium carbonate, sodium carbonate, ammonium carbonate, sodium amino acid carbonate, sodium glycine carbonate, L-lysine carbonate and arginine carbonate.

22. The hydrostatic delivery system according to claim 8, wherein said oxygen precursor is selected from the group consisting of sodium percarbonate, sodium perborate monohydrate, anhydrous sodium perborate, effervescent perborate, and sodium dichloroisocyanurate.

23. The hydrostatic delivery system according to claim 8, wherein said chlorine dioxide precursor is selected from the group consisting of sodium hypochlorite and calcium hypochlorite.

24. The hydrostatic delivery system according to claim 2 wherein the dosage form is a multiparticulate matrix tablet, or capsule.

25. The hydrostatic delivery system according to claim 2 further comprising an enteric coating or one or more pH sensitive barrier polymers.

26. The hydrostatic delivery system according to claim 2, wherein the agent of interest is selected from the group consisting of analgesic, anti-inflammatory, antimicrobial, amoebicidal, trichomonocidal agents, anti-parkinson, anti-malarial, anticonvulsant, anti-depressants, antiarthritics, anti-fungal, antihypertensive, antipyretic, anti-parasite, antihistamine, alpha-adrenargic agonist, alpha blocker, anesthetic, bronchial dilator, biocide, bactericide, bacteriostat, beta adrenergic blocker, calcium channel blocker, cardiovascular drug, contraceptive, decongestants, diuretic, depressant, diagnostic, electrolyte, hypnotic, hormone, hyperglycemic, muscle relaxant, muscle contractant, ophthalmic, parasympathomimetic, psychic energizer, sedative, sympathomimetic, tranquilizer, urinary, vaginal, viricide, vitamin, non-steroidal anti-inflammatory, angiotensin converting enzyme inhibitors, polypeptide, proteins, and sleep inducers.

- 41 -

27. The hydrostatic delivery system of claim 2 further comprising one or more pharmaceutical excipients including but not limited to viscosity enhancer(s), enteric polymer(s), pH-specific barrier polymer(s), diluent(s), anti-adherent(s), glidant(s), binder(s), solubilizer(s), channeling agent(s), wetting agent(s), buffering agent(s), flavorants, adsorbents, sweetening agent(s), colorant(s) and lubricant(s).

28. The hydrostatic delivery system of claim 2 further comprising an adjuvant.

29. The hydrostatic delivery system of claim 2, wherein said hydrostatic delivery system is a matrix-type solid compact, made by a compression or pelletization method.

30. The hydrostatic delivery system of claim 2, wherein said hydrostatic delivery system is a matrix-type extrusion spheroid, made by a wet or dry extrusion method.

31. The hydrostatic delivery system of claim 2, wherein said hydrostatic delivery system is granulated or microencapsulated to form particulates that may be compressed into solid compacts or filled into capsules.

32. The hydrostatic delivery system of claim 2, wherein said hydrostatic delivery system is selected from the group consisting of granulated, particulate, spheroidal, compact, and dry blends, said hydrostatic delivery system can be filled into a capsule or suspended in a suitable liquid vehicle.

1/6




FIGURE 1

2/6




FIGURE 2




FIGURE 3

4/6

Zero-Order Dissolution Profile of Caffeine in Hydrostatic Delivery System
Formulation 10 (Matrix Tablet)




FIGURE 4

Dissolution Profile of RANITIDINE HCL 60 mg Extended Release Tablet




FIGURE 5

6/6

Dissolution Profile Tramadol Hydrochloride 200 mg Extended Release Tablet.




FIGURE 6